

IMMUNE DEFICIENCIES REFERRAL FORM



3100 MacCorkle Avenue S.E. Suite 100
Charleston, WV 25304
TEL: 610-545-6035 | FAX: 610-545-6034
Toll Free: 844-691-5089

Today's Date

- CURRENT PATIENT
 NEW PATIENT

Feb 2018

Patient Name _____ SS# _____ DOB _____
Height _____ Weight _____ Address _____ Apt # _____
 Male Female City _____ State _____ Zip _____
Daytime Tel _____ Cell _____ Email _____
Ship to Patient at Home Work OR Patient will pick up at Physician Office Pharmacy Date Needed _____
Medical History: Cardiac Disease Diabetes Renal Dysfunction IgA Deficient
Allergies _____ Comorbidities _____
Current Medications (if necessary, please fax a complete list) _____

Diagnosis
 D80.0 Hereditary Hypogammaglobulinemia D81.5 Immune Deficiency with Increased Igm
 D83.9 Common Variable Immunodeficiency, unspecified D82.0 Wiskott Aldrich Syndrome
 D81.9 Combined Immunodeficiency, unspecified ICD-10: _____ DX: _____

Insurance Carrier - Primary _____ Name of Insured _____
Relationship _____ ID # _____ Group # _____ Insurance Phone _____
Rx Carrier - Secondary _____ Rx ID # _____ Rx Group # _____ RX Phone _____

Prescriber's Name _____ Office Contact _____
Street Address _____ Suite# _____ City _____ State _____ Zip _____
Tel _____ Fax _____ Email _____
License# _____ NPI# _____ UPIN# _____ DEA# _____

PRESCRIPTION

PLEASE ATTACH COPIES OF PATIENT'S INSURANCE CARDS

Is this the first dose? Yes No If no:
List product _____
Date of last infusion _____
Next dose due _____

ADMINISTER IVIG USING INFUSION PUMP:

- 2 grams/kg over _____ days, as a loading dose, then _____ grams every _____ wk(s) for _____ cycle(s)
 _____ gm/kg or _____ grams every _____ wk(s) for _____ cycle(s)
 Other _____

PRE-MEDICATIONS

- Diphenhydramine (Benadryl) 25-50 mg orally before infusion
 Acetaminophen (Tylenol) 325-650 mg orally before infusion
 Other _____

ADVERSE/ANAPHYLACTIC REACTIONS: PER ELWYN SPECIALTY CARE PROTOCOL

Adults or Children greater than 66 pounds or 30 kg:

- For mild reaction: give Diphenhydramine 50 mg orally, IM or IV and decrease the rate of infusion.
- For moderate reaction: stop infusion, give Diphenhydramine 50mg, orally, IM or IV and contact physician
- For Severe reaction w/breathing problem: stop infusion, call 911, give Epinephrine 0.3mg/0.3ml subcutaneously, Diphenhydramine 50 mg IV or IM. Begin NSS 500ml IV at a rate of 100-150ml/hr and contact physician.

Note: **Dosage adjustment necessary for children less than 30kg or 66 pounds:** Diphenhydramine 1.25mg/kg orally, IM or IV with a maximum of 50mg. If Epinephrine is needed 0.15mg/0.15ml 1:1000 subcutaneously

Nursing: Start PIV as required for administration and nurse to administer infusion in home.

Access: Peripheral PICC Port Other _____

Flushing: Med Center Specialty Pharmacy Protocol (Heparin, 0.9% NaCl, D5W)

Labs _____

By signing this form and utilizing our services, you are authorizing Med Center Specialty Pharmacy and it's employees to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber's Signature (signature required. NO STAMPS) _____ **Date** _____

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately.

Please fax completed referral form to **Med Center Specialty Pharmacy** at **610-545-6034** Visit **WWW.MEDCENTERSPECIALTYPHARMACY.COM** for online fillable forms.